

JC10 Rec'd PCT/PTO 20 DEC 2001

FORM PTO-1390 (Modified)  
(REV 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

112843-039

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

10/019817

INTERNATIONAL APPLICATION NO.  
PCT/EP00/05834

INTERNATIONAL FILING DATE  
23 June 2000

PRIORITY DATE CLAIMED  
30 June 1999

TITLE OF INVENTION

LACTOSE OPERON OF LACTOBACILLUS DELBRUECKII AND ITS USE FOR CONTROLLING GENE  
TRANSCRIPTION AND/OR EXPRESSION IN BACTERIAL CELLS

APPLICANT(S) FOR DO/EO/US

Germond et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
  - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4)
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
10. ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409)
12. ☒ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☒ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

Express Mail No.: EL 727 381 267 US  
Return Receipt Postcard



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Germond et al.  
Appl. No.: PCT/EP00/05834  
Filed: Filed Herewith  
Title: LACTOSE OPERON OF LACTOBACILLUS DELBRUECKII AND ITS USE  
FOR CONTROLLING GENE TRANSCRIPTION AND/OR EXPRESSION IN  
BACTERIAL CELLS  
Art Unit: Unknown  
Examiner: Unknown  
Docket No.: 112843-039

Commissioner for Patents  
Washington, DC 20231

PRELIMINARY AMENDMENT

Sir:

Please amend the above-identified patent application as follows:

**In the Claims:**

Please amend Claims 1, 4, 5-6, 8, and 10-17 as follows:

1. A DNA sequence represented by a general formula selected from the group consisting of:

$p/o - (A)_n - R_y$ , and

$p/o - R_y - (A)_n$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

4. The DNA sequence according to claim 1 wherein the gene coding for a polypeptide of interest is selected from the group consisting of genes encoding enzymes, cell surface proteins, and functional peptides.

5. The DNA sequence according to claim 4, wherein the gene coding for a polypeptide of interest is selected from the group consisting of genes coding for dextransucrase, glycosyltransferase, phytase, transglutaminase, peptidase, phenylalanine ammonia lyase, protease, cell surface antigens, bacteriocins, hormones and insulin.

6. The DNA sequence according to claim 1, which is devoid of catabolite responsive elements.

8. A recombinant microorganism harboring a DNA sequence represented by a formula selected from the group consisting of:

$$p/o - (A)_n - R_y, \text{ and}$$

$$p/o - R_y - (A)_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

10. The microorganism according to claim 8, which is selected from the group consisting of lactic acid bacteria.

11. The microorganism according to claim 8, wherein the DNA sequence is incorporated into the bacteria's chromosome.

12. The microorganism according to claim 8, which is selected from the group consisting of CNCM I-2089, CNCM I-2090 and CNCM I-2091.

13. A method of producing a polypeptide comprising the steps of using a DNA sequence represented by a formula selected from the group consisting of:

$$p/o - (A)_n - R_y, \text{ and}$$

$$p/o - R_y - (A)_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1 for the production of a polypeptide A.

14. The method according to claim 13, wherein the DNA sequence is harbored in a plasmid maintained extra-chromosomal.

15. The method according to claim 13, wherein expression is performed in gram positive microorganisms.

16. The method according to claim 13, wherein expression is performed in microorganisms selected from the group consisting of lactic acid bacteria.

17. A method for the production of food products comprising the steps of using a microorganism having a DNA sequence represented by a formula selected from the group consisting of:

$$p/o - (A)_n - R_y, \text{ and}$$

$$p/o - R_y - (A)_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

Please add newly-submitted Claims 18-19 as follows:

18. The microorganism according to claim 8, wherein the DNA sequence is harbored in a plasmid maintained extra-chromosomal.

19. The method of claim 13, wherein the DNA sequence is in the bacteria's chromosome.

#### REMARKS

This Preliminary Amendment is submitted in the above-identified patent application. Pursuant to the Preliminary Amendment, Claims 1, 4, 5-6, 8, and 10-17 have been amended and newly-submitted Claims 18-19 have been added. This Preliminary Amendment does not add new subject matter. Applicants also note for the record that the purpose of this Preliminary Amendment is to place the claims in proper format and/or add additional claims and not for purposes of narrowing the claims. This Preliminary Amendment is not being submitted for purposes of patentability and therefore Applicants do not intend to disclaim any subject matter in view of this Preliminary Amendment.

Attached hereto is a marked-up version of the changes made to the specification and claims by the current amendment. The attached page is captioned "**Versions with Markings to Show Changes Made.**"

Respectfully submitted,

BELL, BOYD & LLOYD LLC

BY 

Robert M. Barrett  
Reg. No. 30,142  
P.O. Box 1135  
Chicago, Illinois 60690-1135  
Phone: (312) 807-4204

**VERSION WITH MARKINGS TO SHOW CHANGES MADE**

**In the Claims:**

Please amend Claims 1, 4, 5-6, 8, and 10-17 as follows:

1. (Amended) A DNA sequence represented by a [the following] general formula  
selected from the group consisting of:

p/o - (A)<sub>n</sub> - R<sub>y</sub>, [or] and

p/o - R<sub>y</sub> - (A)<sub>n</sub>

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to  
bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

4. (Amended) The DNA sequence according to claim 1 wherein the gene coding  
for a polypeptide of interest is selected from the group consisting of genes encoding enzymes,  
cell surface proteins, [or] and functional peptides.

5. (Amended) The DNA sequence according to claim 4, wherein the gene coding  
for a polypeptide of interest is selected from the group consisting of genes coding for  
dextranucrase, glycosyltransferase, phytase, transglutaminase, peptidase, phenylalanine  
ammonia lyase, protease, cell surface antigens, bacteriocins, hormones [or] and insulin.



6. (Amended) The DNA sequence according to [any of the preceding claims] claim 1, which is devoid of catabolite responsive elements.

8. (Amended) A recombinant microorganism harboring a DNA sequence [according to any of the preceding claims] represented by a formula selected from the group consisting of:

$$\text{p/o} - (\text{A})_n - \text{R}_y, \text{ and}$$

$$\text{p/o} - \text{R}_y - (\text{A})_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of Lactobacillus delbrueckii;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

10. (Amended) The microorganism according to claim 8 [or 9], which is selected from the group consisting of lactic acid bacteria.

11. (Amended) The microorganism according to [any of the claims] claim 8 [to 10], wherein the DNA sequence [of claims 1 to 7] is incorporated into the bacteria's chromosome [or is harbored in a plasmid maintained extra-chromosomal].

12. (Amended) The microorganism according to claim 8, which is selected from the group consisting of CNCM I-2089, CNCM I-2090 [or] and CNCM I-2091.

13. (Amended) [Use of] A method of producing a polypeptide comprising the steps of using a DNA sequence [according to any of the claims 1 to 7] represented by a formula selected from the group consisting of:

$$\text{p/o} - (\text{A})_n - \text{R}_y, \text{ and}$$

$$\text{p/o} - \text{R}_y - (\text{A})_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of Lactobacillus delbrueckii;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1 for the production of a polypeptide A.

14. (Amended) The [use] method according to claim 13, wherein the DNA sequence is harbored in a plasmid maintained extra-chromosomal [or is in the bacteria's chromosome].

15. (Amended) The [use] method according to [any of the claims] claim 13 [or 14], wherein expression is performed in gram positive microorganisms.

16. (Amended) The [use] method according to [any of the claims] claim 13 [to 15], wherein expression is performed in microorganisms selected from the group consisting of lactic acid bacteria.

17. (Amended) [The use of a microorganism according to any of the claims 8 to 11] A method for the production of food products comprising the steps of using a microorganism having a DNA sequence represented by a formula selected from the group consisting of:

$$\text{p/o} - (\text{A})_n - \text{R}_y, \text{ and}$$

$$\text{p/o} - \text{R}_y - (\text{A})_n$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9, which retains its capability to bind to the lac repressor protein of Lactobacillus delbrueckii;

A denotes a heterologous gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2; and

Y is 0 or 1.

Please add newly-submitted Claims 18-19 as follows:

18. The microorganism according to claim 8, wherein the DNA sequence is harbored in a plasmid maintained extra-chromosomal.

19. The method of claim 13, wherein the DNA sequence is in the bacteria's chromosome.

Rec'd PCT/PTO 20 DEC 2001

WO 01/02576

10/019817  
PCT/EP00/05834

1

**The lactose operon of Lactobacillus delbrueckii and its use for  
controlling gene transcription and/or expression in bacterial cells**

The present invention relates to a DNA sequence suitable for controlled transcription and/or expression of a variety of different genes in bacteria, preferably in gram positive bacteria. In particular, the present invention pertains to a DNA sequence comprising the promoter/operator region of the lac operon of Lactobacillus delbrueckii with a DNA sequence coding for a gene product of interest being arranged in functional relationship therewith.

In the food industry a variety of different lactic acid bacteria are used to prepare food products, such as cheese, buttermilk or yogurt, essentially relying on the fermentative activity of said bacteria. For this purpose bacteria of e.g. the genus Streptococcus or Lactobacillus are put to use, with their primary function being to convert sugar, such as lactose or glucose, to lactic acid and to produce a satisfying texture and aroma.

A problem of any such fermentation resides in that the end product of the metabolic process, lactic acid, is often produced abundantly rendering the dairy product acid to an extent not accepted by most of the consumers. Thus, the persons attending the production process must take care so as to control the progress of the manufacturing process in order to avoid any detrimental effects of an excessive fermentation to the food product.

Lactic acid bacteria have also been used as fermenting agents for the preservation of food taking benefit of a low pH and the action of fermentation products generated during the fermentation process to inhibit the growth of spoilage bacteria. Also in such cases it is up to the personal to control the biological activity of the micro-organisms and to bring the production process at a desired stage to a stop.

Therefore, there is a need in the art to control the fermentative activity of micro-organisms used for the production of food, thus being capable to provide a fermentation up to a desired stage of the production process of the food product without involving too much labor work from the side of the attending staff.

In the recent past producers have tried to add valuable properties to bacteria utilized in fermentative processes via recombinant technologies. To this end, a vector bearing a homologous or heterologous gene coding for a polypeptide of interest is brought into the bacterial cell to either remain extra-chromosomal or to be inserted in the bacteria's chromosome. Expression is then effected by means of a heterologous promoter linked to the respective gene, which may be controlled as desired.

Though a control of the expression of the heterologous gene may be achieved thereby, this procedure is disadvantageous in that the regulon used to control the expression of the desired gene is in most cases exogeneous to the bacteria. Yet, the use of foreign gene material in bacteria utilized for the preparation of food products is presently not accepted by the consumer.

An object of the present invention is to provide means for a controlled transcription/expression of genes in bacteria that do not exhibit the disadvantages of the prior art techniques.

The above object has been solved by providing a DNA sequence exhibiting the following general formula

$$\begin{aligned} p/o - (A)_n - R_y, \text{ or} \\ p/o - R_y - (A)_n \end{aligned}$$

wherein p/o denotes the DNA sequence identified under SEQ ID No. 9 or a functional variant thereof, which retains the capability to bind the lac repressor protein of *Lactobacillus delbrueckii*; A denotes a gene coding for a polypeptide of interest; n denotes an

integer of  $\geq 0$ ; R denotes the gene coding for the lacZ repressor protein as identified under SEQ ID No. 2 or a functional variant thereof; and Y is 0 or 1.

Consequently, in case  $y = 0$  the gene coding for the lac repressor protein may be situated at any site of the bacteria's chromosome and may be subject to any suitable promoter region such that is e.g. expressed constitutively.

According to a preferred embodiment the DNA sequence is represented by the general formula  $p/o - (A)_n - R$  or  $p/o - R - (A)_n$ , respectively, with p, A and R being as identified above. In this case, i.e. in the case of  $Y = 1$ , the construct is composed of three components, the promoter/operator region, the gene coding for the polypeptide of interest and the gene coding for the lac repressor protein (lacR), with the genes coding for the respective polypeptides (A, lacR) being arranged in either order. The lacR gene is such subject to its own promoter/operator. In addition, the gene coding for the lacR protein may be arranged such that its transcriptional direction is opposite to that of the gene A coding for the polypeptide of interest.

According to another preferred embodiment the gene coding for a polypeptide of interest is selected from group consisting of genes encoding enzymes, cell surface proteins or functional peptides, such as e.g. dextransucrase, glycosyltransferase, phytase, transglutaminase, peptidase, phenylalanine ammonia lyase, protease, cell surface antigens, bacteriocins, hormones, insulin.

The promoter region may also lack any catabolite responsive elements, so that no repression in the presence of a particular carbon source may be effected.

According to another preferred embodiment the DNA sequence is used to transform a microorganism, which is preferably a gram positive bacterium or more preferably a microorganism selected from the group consisting of lactic acid bacteria, such as Lactococcus, Lactobacillus, Streptococcus, Leuconostoc etc..

The DNA sequence may be incorporated into the respective microorganism by means of an appropriate vector, such as a plasmid or a viral construct, and may remain in the microorganism extra-chromosomal. Yet, according to another preferred embodiment the construct may also be inserted into the bacteria's chromosome by means of conjugation followed by a crossing over between the plasmid and a homologous chromosomal sequence, or direct transformation and single or double cross over integration.

The DNA construct according to the present invention may therefore suitably be used for the transcription and/or expression of gene products, such as RNA or polypeptides.

In the figures,

Fig. 1 shows a comparison of *L. delbrueckii* promoter sequences;

Fig. 2 shows the organisation of the promoter region of *L. delbrueckii* LL44 and LB68;

Fig. 3 shows the organisation of the promoter region of *L. delbrueckii* N299;

Fig. 4 shows the nucleotide and amino acid sequence of the *L. delbrueckii* subsp. *lactis* LL44 *lacR* gene;

Fig. 5 shows a physical map of the lactose operon of different *L. delbrueckii* strains;

Fig. 6 shows a schematic representation of the construction of pLL 62;

Fig. 7 shows a schematic representation of the construction of pLL110 and pLL112 (CNCM I-2089);

Fig. 8 shows a schematic representation of the construction of pLL113, pLL115 (CNCM I-2090) and pLL116 (CNCM I-2091);

Fig. 9 shows the results of the expression of  $\beta$ -glucuronidase using the constructs of the present invention;

Fig. 10 shows the results of experiments, wherein constructs of the present invention were used to express  $\beta$ -galactosidase in *L. delbrueckii*.

The correct regulation of gene transcription and/or translation is a prerequisite for any cellular system to maintain its metabolic and catabolic turn-over and to respond to changes of the environment. In order to better adopt to environmental changes, such as a change of the nutritional source, microorganisms have developed various systems to control gene expression on both, the transcriptional and/or the translational level.

For the gram negative bacterium *Escherichia coli* several of such control systems have been studied. To this end, it has been found that the genes responsible for the catabolism of lactose, that is the genes coding for  $\beta$ -galactosidase (*lacZ*), lactose permease (*lacY*) and transacetylase (*lacA*), respectively, are organized in a particular structural pattern which was termed operon. In this operon the genes coding for the structural polypeptides are located downstream a DNA region coordinating the transcription of said genes (operator) which region is negatively controlled by a repressor protein. The gene coding for the repressor protein is in turn controlled by a different promoter, which allows for a constitutive expression of the repressor.

Under repressive conditions the repressor protein binds to the promoter/operator region thus preventing transcription of the adjacent genes (*lacZYA*). Yet, in the presence of lactose or allolactose, which represent inducing compounds, the repressor forms a complex with the inducer, which complex has only a limited affinity to the promoter/operator, thus giving way for a transcription of the genes.

The regulation of the *lac* operon in gram positive bacteria, such as lactic acid bacteria, has been found to be also mediated via a polypeptide acting as a repressor, which is encoded by a gene termed *lacR*. This gene has been sequenced for *Lactococcus lactis*



(van Rooijen et al., J. Biol. Chemistry **265** (1990), 18499 - 18503) and *Staphylococcus aureus* (Oskouian et al., J. Bacteriol. **169** (1987), 5459 - 5465). Also in these microorganisms the repressor has a high affinity for the operator and binds to it in the absence of an inducer, resulting in a transcriptional blockade of genes located downstream thereof. In the presence of an inducer molecule, such as lactose or galactose, the repressor is not capable to bind to the operon any more with the all structural genes following the operator region being transcribed together.

In lactic acid bacteria two different systems for the internalisation of lactose into the cells have been reported. Poolman et al. described in J. Bacteriol. **171** (1989), 244-253 a phosphoenolpyruvate dependent phosphotransferase system and a lactose permease system. It is reported that in *Lactobacillus*, *Streptococcus* and *Leuconostoc* lactose permease acts as a lactose/galactose antiporter system. The microorganisms internalize lactose from the medium as free sugar which is subsequently hydrolyzed to glucose and galactose. Galactose is either converted by the Leloir pathway into glucose-6-phosphate or is released into the medium by the permease.

In J. Bacteriol. **171** (1989), 244-253 and **173** (1991), 1951 - 1957 the cloning and sequencing of the lactose permease genes (*lacS*) of *S. thermophilus* and *L. delbrueckii*, respectively, have been reported. The genes were found to be arranged in an operon structure each together with a gene coding for  $\beta$ -galactosidase (*lacZ*). The gene for  $\beta$ -galactosidase was disclosed for several lactic acid bacteria, such as *S. thermophilus* (Schroeder et al., J. Gen. Microbiol. **137** (1991), 369-380), *L. bulgaricus* (Schmidt et al., J. Bacteriol. **171** (1989), 625-635) and *L. casei*. (Chassy et al., FEMS Microbiol. Rev. **63** (1989), 157-166).

Yet, it has been found that the *lac* genes may also be subject to catabolite repression. In the presence of particular compounds, such as e.g. glucose, the genes of the *lac* operon are maintained in a repressed condition. This action is mediated by a cis acting element, that has been first described in *Bacillus subtilis* (Weickert et al., Proc. Natl. Acad. Sci. USA **87** (1990), 6238 - 6242) and has been termed "Catabolite Responsive Element"

(CRE). This element is present in a variety of gene arrangements encoding carbon catabolite enzymes in different gram positive microorganisms and seems to be controlled by the trans acting factors CcpA (Catabolite control protein A) and Hpr. In the presence of glucose both of these proteins bind to the CRE sequence acting as a negative regulation of transcription.

The species *L. delbrueckii* contains mainly two subspecies, *L. lactis* and *L. bulgaricus*, which are defined by several physiological and genetic criteria. One of these criteria concerns the regulation of the lactose (lac) operon expression. In the subspecies *lactis*, the expression is induced by lactose, that is the lac genes are expressed only in the presence of this sugar. In the subspecies *bulgaricus*, which has been selected a long time ago for its ability to ferment milk for yogurt production, the control of the lac genes expression was lost probably due to the constant availability of lactose in milk and the genes are constitutively expressed in the presence or in the absence of lactose.

For the experiments leading to the present invention six representative strains have been chosen to study the regulation of genes via the lac operon. Five strains belong to the subspecies *lactis* and have been termed LL44, LB68, N62 and N141, which are inducible by the presence of lactose, and LB10, that has been classified as belonging to the subspecies *lactis* but with a constitutive expression of the lac operon. The sixth strain, N299, corresponding to ATCC 11842, was chosen as a typical *bulgaricus* strain, with a constitutive expression of the lac genes.

During the analysis of the lactobacillus strains the region flanking the lactose (lac) operon was isolated and the following regions could be characterized.

### **The lac A gene**

During the study of the promoter region of the lac operon, an open reading frame (ORF) of 570 bp was discovered upstream the lacS gene in the same orientation. This gene was completely sequenced for the *L. delbrueckii* subsp. *lactis* strain LL44 (SEQ ID. No 1) and it encodes a polypeptide of 190 amino acid residues (SEQ ID. No. 2). The presence of a

putative promoter and a potential  $\rho$ -independent-termination signal (stem-loop structure ; position 823, SEQ. ID. NO 1) suggest that this gene does not belong to the lac operon.

### **The promoter region**

The region comprised between the end of lacA and the beginning of lacS (lactose permease) was sequenced and showed several attributes specific for a promoter. Two palindromic sequences, called O1 and O2 (Fig. 1), were discovered and may serve as operators for the binding of the lac repressor (LacR). The operators of the 6 strains studied (LL44, LB68, N62, N141, LB10 and N299) were sequenced. Several differences were found, which were partly due to the insertion of IS-elements in the O1 inverted repeats and partly due to small nucleotides changes in the sequence (Fig. 1). ISL7 is inserted in N141 exactly at the TGT motif of the 5' end of O1, but restoring the bases TG (Fig. 1). In N299, the *L. delbrueckii* subsp. *bulgaricus* strain, the insertion of ISL5 at the same position destroyed the TGT motif, which might explain the constitutive transcription of adjacent genes.

In lactic acid bacteria effective repression of transcription is deemed to involve the cooperative binding of LacR to two different operator sequences. In *L. delbrueckii*, a small operator, called O2, was found 4 bp downstream O1. The length of O2 showed considerable variation in length between the 6 strains analysed due to small nucleotide changes in the sequence (Fig. 1). However, no IS-element was found in this operator. The core of both inverted repeats is composed of the nucleotides TGTTTA (SEQ. ID. No. 3) and (SEQ. ID. No. 4), except in LL44 where the final A was replaced by G in O2 (SEQ. ID. No. 5).

A sequence of 14 nucleotides homologous to the catabolite responsive element (CRE) (Weickert, supra), was discovered 40 nucleotides upstream the lacS start codon of all the *L. delbrueckii* strains studied (Fig. 2, Fig. 3, SEQ ID No. 8)). Catabolite repression is effective in many bacteria and acts at the level of transcription via negative transcriptional control. It involves a cis-acting element mediating repression of the genes under its control in the presence of glucose. The sequence of these elements is highly conserved among

different species and the one of *L. delbrueckii* subsp. *lactis* is highly similar to other elements.

The figures 2 and 3 present a view of the promoter region of the lac operon of 5 of the strains studied. They are considered as representative of the different possible lac operon organisations in the *L. delbrueckii* species. Four inducible *lactis* strains, LL44, LB68, N62 and N141 and one constitutive *bulgaricus* strain, N299 were represented. Strain LB10 was not shown, because its promoter region is identical to N62 except that ISL6 is not present in the lac promoter of this strain. The initiation of transcription in N299 (Leong-Morgenthaler et al., J. Bacteriol. 173 (1991), 1951-1957) falls in the middle of the CRE element, just after the three nucleotides change that were found in the CRE element of this strain (Figs. 2 and 3).

LL44 was chosen as a reference due to the absence of any IS-element in its lac operon region. In this strain, the lacA gene is followed by the two operators O1 and O2 and the CRE element upstream the lacS gene. In other strains, the presence of IS-elements changed the promoter sequence and particularly the O1 operator sequence.

### The lac R gene

Surprisingly, and in contrast to the gene arrangements of known lac operons the gene coding for the repressor protein was discovered downstream the lacZ gene in *L. delbrueckii* subsp. *lactis* LL44 and LB68, with the result that the repressor is under its own transcriptional control.

The region covering the lacR gene was PCR amplified using the following primers:

CGCCTGGTGATTCAGCC (SEQ ID No. 6)

AGCTTTACGGGGAAGTCGGG (SEQ ID No. 7)

which are located at the end of the  $\beta$ -galactosidase (lacZ) gene (SEQ ID No. 6) and in the Asn-tRNA synthetase (asnA) gene (SEQ ID No. 7).

Sequencing of this region revealed the sequence identified under SEQ ID No. 8, which exhibits an open reading frame of 999 bp in the same orientation as lacZ. The lacR gene is preceded by a ribosome binding site (RBS) and followed by a putative  $\rho$ -independent-termination signal (stem-loop structure ; position 1149). The putative amino polypeptide derived therefrom is shown under SEQ ID No. 2 (333 amino acid residues).

Computer-assisted analysis of the *L. delbrueckii* lacR gene predicted the protein secondary structure of the beginning of the gene (positions 4 to 23) to be a helix-turn-helix motif, which represents the highest homology region with other repressors. This type of protein secondary structure is a common feature for DNA-binding proteins and binds to the operators located in the promoter region.

### The lac operon

The organisation of the lac operon of different *L. delbrueckii* strains is represented in figure 5. The lac operon per se is preceded by the thiogalactosyl-transacetylase (lacA) gene, which is followed by a rho-independent termination signal. This gene is followed by the promoter region of the lac operon, which comprises the inverted repeats (operators) involved in the regulation of the gene expression and the CRE sequence. The lac operon is composed of three genes, the permease (lacS), the  $\beta$ -galactosidase (lacZ) and the repressor (lacR). The lacS and lacZ genes are separated by 4 nucleotides, whereas lacR is distant from lacZ by 52 nucleotides.

Five IS-elements have been found in the promoter and the terminator region of the lac operon, which were called ISL3, ISL4, ISL5, ISL6 and ISL7. One of these Elements, ISL3, has been described in Germond et al, Mol. Gen. Genet. **248** (1995), 407-416.

The lactose permease (lacS), the  $\beta$ -galactosidase (lacZ) and the repressor (lacR) constitute the lac operon per se. The three genes are linked together as a lacSZR operon in *L. delbrueckii* subsp. *lactis* without any promoter in between. This gene arrangement is not a common feature for regulated operons, as the lacR gene is normally not a part of the operon.

Induction of  $\beta$ -glucuronidase production showed considerable differences between the different promoters associated to LL44 lacR. The plasmid carrying the LL44 promoter (pLL112) was fully induced by 1.0% lactose up to levels of 250 mU /mg protein, whereas plasmids carrying the N299 promoter (pLL116) reached about 100 mU/mg protein when the lacR gene was in the reverse orientation as the gusA gene. When the lacR gene was

introduced in the same orientation as the gusA gene (pLL115), levels of induction dropped to around 40 mU/mg protein. This may be explained by the absence of a promoter in front of lacR and its control either by the N299 strong lac promoter when it is in the sense of the gusA gene or by a weak promoter which must be located on pNZ272 when it is orientated antisense. It was also not possible to obtain the lacR gene in the same sense than gusA with the LL44 promoter. Production of GusA also showed that all plasmids were subjected to glucose catabolite repression due to the presence of a CRE sequence in the promoter of *L. delbrueckii*. The CRE of N299 is partly disrupted and the repression observed with this promoter is weaker than with LL44.

Measurements of the  $\beta$ -galactosidase activity in vivo revealed that the *L. delbrueckii* strains were almost not subjected to glucose catabolite repression. The  $\beta$ -galactosidase activity also confirmed the inducible characteristics of the *L. delbrueckii* subsp. *lactis* strains and the constitutivity of the *bulgaricus* strains. Indeed, LL44, LB68, N62 and N141 were not induced in the presence of glucose as sole carbon source, whereas in N299  $\beta$ -galactosidase activity was also high in the glucose medium. The inducible characteristic of an operon allows the strain to quickly switch from metabolizing one substrate to another.

Thus, according to the present invention the transcription and/or expression of homologous and/or heterologous genes in bacteria may be suitably obtained by preparing the construct p/o-(A)<sub>n</sub>-R<sub>y</sub> or p/o-R<sub>y</sub>-(A)<sub>n</sub> as illustrated above and introducing said construct into the bacterium.

According to the present invention it is now possible to control the transcription and/or expression of a specific gene polypeptide by adjusting the content of glucose and/or lactose, respectively, in the culture medium. As an example, the bacterium may be grown in a medium lacking lactose to a desired degree and in a next step may then be transferred to a medium containing lactose or lactose is simply added to the medium. Only at this stage the repressive activity of the construct on the transcription and/or expression of the gene of interest will be relieved and the gene product will be produced.

In order to assist the inductive activity of the construct the gene for the lactose permease may be one of the polypeptides designated "A". To this end, if an amount of lactose enters the bacterial cell the genes following the promoter described here are transcribed with one of the polypeptides to be expressed thereby being lactose permease, which actively transfers additional lactose into the cell.

Further, it may well be suitable to delete the region harboring the catabolite responsive element, so that the construct is not subject to carbon repression.

The following examples illustrate the invention without limiting the scope of the appended claims.

### **Example 1**

#### **Bacterial strains and growth conditions**

Lactobacilli were grown in Difco Lactobacillus MRS-broth (DeMan et al., J. Appl. Bacteriol. **23** (1960), 130-135) or in Oxoid BHI-broth. Lactococci strains were cultivated in Difco M17-broth (Terzaghi et al., Appl. Microbiol. **29** (1975), 807-813) supplemented with 0.5% glucose (GM17). E. coli was grown in YT-broth 2x prepared according to Maniatis et al., Molecular cloning : A Laboratory Manual, 2<sup>nd</sup> ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. (1989). All strains were stored at -80°C in their respective media containing 10% glycerol ten times concentrated.

For plasmid selection and maintenance in Escherichia coli, ampicillin (100 µg/ml), chloramphenicol (30 µg/ml), or kanamycin (50 µg/ml) were added to the growth medium (ampicillin and kanamycin obtained from Sigma, chloramphenicol from Boehringer Mannheim). For plasmid selection and maintenance in Lactococcus lactis MG1363, chloramphenicol at 12 µg/ml was added to the growth medium.

#### **Chromosomal and plasmid DNA purification**

Lactobacilli DNA was prepared according to Delley et al., Appl. Environ. Microbiol. **56** (1990), 1967-1970. Plasmid DNA from E. coli was purified by the alkali lysis method (Maniatis, supra). Lactococcus plasmid DNA was prepared following the same procedure,



except that the cultures were first incubated 30 min at 37°C in TE (10 mM Tris-HCl, pH 8.0 ; 1 mM EDTA) containing 10 mg/ml lysozyme (all chemicals obtained from SIGMA).

#### DNA manipulations and bacterial transformations

Agarose gel electrophoresis, restriction enzyme digestions, ligations and transformations in *E. coli* were performed according to standard procedures (Maniatis, supra). When necessary, the digested DNA was blunt-ended with T<sub>4</sub> DNA polymerase (Boehringer-Mannheim) directly in the digestion mixture for 5 min at 37°C.

The different *L. delbrueckii* operators were cloned into plasmid pKK232-8 (Ap<sup>r</sup>, Cam<sup>r</sup>, contains a promoterless chloramphenicol acetyltransferase (cat) gene, *E. coli* vector, Pharmacia) and transformed in *E. coli* SURE (obtained from STRATAGENE). Transformants were selected on YT-agar plates containing 30 µg/ml chloramphenicol. The LL44 lacR gene was cloned into plasmid pKK223-3 (Ap<sup>r</sup>, contains a strong tac promoter, Pharmacia) and transformed in *E. coli* SURE. Transformants were selected on YT-agar plates containing 100 µg/ml ampicillin. For lacR overexpression, 2 mM isopropyl-β-D-thiogalactoside (IPTG) was added to the culture and cells were allowed to grow for an additional 4-5 hours at 37°C. Cultures were centrifuged and the pellets were resuspended in 0.1 M Hepes, sonicated and centrifuged. Supernatants were applied to a SDS/PAGE gel (Laemmli, Nature 227 (1970), 680-685).

When the lacR gene was cloned into plasmid pET11c (obtained from STRATAGENE), plasmids were transformed in *E. coli* BL21 (STRATAGENE). Transformants were selected on YT-agar plates containing 100 µg/ml ampicillin, and lacR was overexpressed as described above in the presence of 1 mM IPTG. The lacR gene was finally cloned into plasmid pACYC177 (New England Biolabs) into the PstI site of the ampicillin gene and transformed in *E. coli* XL1-Blue (STRATAGENE). Transformants were selected on YT-agar plates containing 50 µg/ml kanamycin. Plasmid pACYC177 containing the lacR gene was cotransformed with the pKK232-8 plasmids containing the different operators in *E. coli* XL1-Blue. Transformants were selected on 100 µg/ml ampicillin for the presence of pKK232-8 and 50 µg/ml of kanamycin for the presence of pACYC177.

Competent cells and transformations of *Lactococcus lactis* MG 1363 (plasmid free derivative of NCDO 712; Gasson, M.J., NCDO (1983) 712) and other lactic streptococci after protoplast-induced curing, J. Bacteriol. 154:1-9) were realized according to Holo et al. Appl. Environ. Microbiol. 55 (1989), 3119-3123. Cells were grown in GM17-broth containing 500 mM sucrose and 3% glycine to an OD<sub>600</sub> of 0.2 - 0.3, centrifuged and washed several times in a solution containing 500 mM sucrose and 10% glycerol. Competent cells were stored 100x concentrated at -80°C in the same solution until use.

Cells (40 µl) were mixed with 5 µl of plasmid DNA and electrotransformed at 2.0 kV, 25 µF and 200 Ω. Cells were resuspended in GM17-broth containing 20 mM MgCl<sub>2</sub> and 2 mM CaCl<sub>2</sub> and expressed for 90 min at 30°C. Cells were then plated on GM17-agar containing 12 µg/ml of chloramphenicol. When the different *L. delbrueckii* promoters were cloned upstream the GUS gene, 50 µg/ml of X-glu (5-bromo-4-chloro-3-indolyl-β-D-glucuronide) (Amresco) was added to the GM17-agar plates to select for blue colonies.

#### DNA amplification and sequencing

DNA amplification was done by polymerase chain reaction (PCR) in the presence of 0.2 mM of each dNTPs, 1 µM of each oligonucleotide primers, the appropriate buffer and the Taq polymerase (Gold polymerase, Perkin-Elmer) under the following conditions : 1 min at 94°C - 2 min at 55°C and 3 min 30 at 72°C for 35 cycles. Custom made oligonucleotide primers were used. DNA sequences were determined directly by PCR using the VISTRA Thermo sequenase sequencing kit (Amersham) (Sanger et al., Proc. Natl. Acad. Sci. USA 74 (1977), 5463-5467) and then analysed with the help of the University of Wisconsin, USA, Genetics Computer Group (GCG) computer software package (Devereux, Nucleic Acids. Res. 12 (1984), 387-395).

#### Isolation of spontaneous repressor negative LL44 mutants

The *L. delbrueckii* subsp. *lactis* strain LL44 was isolated on MRS-agar containing glucose and 200 µg/ml X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside) (Sigma) to check the inducible state of its lac operon. One white colony was further incubated overnight at 42°C in MRS-broth containing 10% lactose. The culture was stored refrigerated for 24 hours and then subcultured in MRS-broth containing 10% lactose for 7h

at 42°C. This culture was diluted and plated on Oxoid Reinforced Clostridial Medium (RCM) agar containing only glucose and 200 µg/ml X-gal to select for a blue lac constitutive mutant. Plates were incubated for 72 h at 40°C under microaerophilic conditions (BBL, microaerophilic system No. 71034). A blue colony was obtained, further cultivated and stored in MRS-broth as described above.

Chloramphenicol acetyltransferase assay (Shaw, Methods Enzymol. 43 (1975), 737-755.

Cells were grown overnight in YT-broth 2x containing the necessary antibiotics (see DNA manipulations and bacterial transformations) at 37°C under shaking. 10 ml of fresh YT-broth containing either 0.5% lactose or 0.5% glucose was inoculated with 1% of the overnight culture. Kanamycin (50µg/ml) and/or ampicillin (100 µg/ml) were used as antibiotics. Cells were incubated for 3 hours at 37°C, centrifuged and the pellet was resuspended in 2 ml Hepes 0.1 M. The suspension was sonicated for 30 sec, spun down and the supernatant was mixed with 1 volume of buffer 2x concentrated (200 mM Tris-HCl, [pH 8.0] – 0.2 mM Acetyl CoA – 0.8 mg/ml DTNB) and chloramphenicol was added at a final concentration of 0.1 mM. The mixture was incubated at 37°C for 4 min and the OD red at 412 nm.

In a second cycle of experiments, the YT-broth was replaced by a solution containing : 1/5 vol M9-salts 5x (Na<sub>2</sub>HPO<sub>4</sub> • 7H<sub>2</sub>O, 450 mM – KH<sub>2</sub>PO<sub>4</sub>, 110 mM – NaCl, 45 mM – NH<sub>4</sub>Cl, 93 mM), 0.5 % glucose or lactose, 1 mM CaCl<sub>2</sub>, 20 mM MgCl<sub>2</sub> and 10 mM thiamine. The rest of the assay was realised as described above.

Determination of β-glucuronidase activity

(modified from Bergmeyer, Methods of Enzymatic Analysis 2 1983, 206-209)

Starters of transformed *Lactococcus lactis* were prepared by incubating the bacteria overnight at 30°C in M17-broth containing 0.5% mannose and 12 µg/ml chloramphenicol. Ten ml of M17 containing different sugars were inoculated at 2% with the overnight culture and incubated at 30°C to an OD<sub>660</sub> of about 1.0. Following centrifugation, cells were stored at -20°C until use. For the assay, they were resuspended in 2 ml of distilled water containing 0.1% Triton X100, incubated 30 min at room temperature and collected by centrifugation. Cells were then resuspended in 2 ml of GUS buffer (50mM NaHPO<sub>4</sub> [pH 7.0], 10 mM β-mercaptoethanol, 1 mM EDTA, 0.1% Triton X100) containing 50

$\mu\text{g/ml}$  of para-nitrophenyl- $\beta$ -D-glucuronide (Clonotech). After 1 hour incubation at  $37^\circ\text{C}$ ,  $500\ \mu\text{l}$  were mixed with  $500\ \mu\text{l}$  of stop solution ( $2\ \text{mM Na}_2\text{CO}_3$ ). Centrifugation was followed by absorbance measurement at  $415\ \text{nm}$ .  $\beta$ -glucuronidase activity was calculated by using the equation :

$$\text{U/mg prot.} = \text{OD}_{415} \times \text{dilution factor} / 18 \times t \times \text{mg prot./ml}$$

wherein  $t$  is the reaction time, 18 the millimolar extinction coefficient of the para-nitrophenol and 0.4 the dilution factor.

The protein content of the culture was determined as described by Bradford. Anal. Biochem. **12** (1976), 248-254, using the Bio-Rad protein assay with bovine serum albumin as standard. Samples of  $0.1\ \text{ml}$  of the culture were mixed with  $5\ \text{ml}$  Bio-Rad dye reagent (diluted 1:4 with distilled water). After incubation for 5 min, the absorbance was read at  $550\ \text{nm}$ .

For the  $\beta$ -glucuronidase induction, fresh M17-broth containing 0.5% mannose and  $12\ \mu\text{g/ml}$  chloramphenicol was inoculated at 2% with the starter cultures and incubated at  $30^\circ\text{C}$  to an  $\text{OD}_{660}$  of about 0.2. The cultures were split and the different sugars added. Cultures were further grown at  $30^\circ\text{C}$  until the  $\text{OD}_{660}$  reached 1.0.  $\beta$ -glucuronidase activity was measured as described above at 60 min intervals.

#### Determination of $\beta$ -galactosidase activity (modified from Bergmeyer, supra)

Starters of *L. delbrueckii* were prepared by incubating bacteria overnight in MRS-broth at  $42^\circ\text{C}$ .  $10\ \text{ml}$  of BHI containing different sugars (table I) were inoculated at 2% with the overnight culture and incubated at  $42^\circ\text{C}$  to an  $\text{OD}_{600}$  of about 1.0. For the *Lactobacillus delbrueckii* subsp. *lactis* strain N141, galactose was replaced by mannose (as galactose induced the lac operon). Following centrifugation, cells were stored at  $-20^\circ\text{C}$  until use. For the assay, pellets were resuspended in  $2\ \text{ml}$  of distilled water containing 0.1% Triton X100, incubated at room temperature for 30 min, and collected by centrifugation. Cells were resuspended in  $2\ \text{ml}$  of Z buffer ( $0.06\ \text{M Na}_2\text{PO}_4$ ;  $0.04\ \text{M NaH}_2\text{PO}_4 \cdot 7\text{H}_2\text{O}$ ;  $0.01\ \text{M KCl}$ ;  $0.001\ \text{M MgSO}_4 \cdot 7\text{H}_2\text{O}$ ;  $0.05\ \text{M } \beta$ -mercaptoethanol [pH 7.0]) (Miller, Experiments in Molecular Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1972),

diluted 10 times in the same buffer containing 50 µg/ml ortho-nitrophenyl-β-D-galactopyranoside (ONPG). After 30 min incubation at 42°C, 200 µl were mixed with 800 µl of stop solution (400 mM Na<sub>2</sub>CO<sub>3</sub>; 50 mM EDTA) Centrifugation was followed by absorbance measurement at 420 nm. β-galactosidase activity was calculated by using the equation :

$$U/OD_{600} = OD_{420} \times \text{dilution factor} / 3.5 \times t \times OD_{600},$$

wherein t is the reaction time, 3.5 the millimolar extinction coefficient of the ortho-nitrophenol and 10 the dilution factor.

For the induction of β-galactosidase, fresh BHI-broth containing 0.5% galactose was inoculated with the starter cultures and incubated at 42°C to an OD<sub>600</sub> of about 0.2. The cultures were split and the different sugars were added. Cultures were further grown at 42°C to an OD<sub>600</sub> of about 1.0. β-galactosidase activity was measured as described above at 15 min intervals.

## Example 2

### Expression of the L. delbrueckii LL44 lacR gene in E. coli

The lac repressor (lacR) gene of L. delbrueckii strain LL44 was tentatively expressed in E. coli. It was first PCR amplified using the following primers

Pst I  
ATATTACTGCAGAGTAAAAGCGAGT SEQ ID No. 10

Hind III  
ATAAATAAGCTTACAGAATGCAGCC SEQ ID No. 11

which contain a PstI and a HindIII site, respectively. The fragment was cloned in the corresponding sites behind the strong tac promoter of pKK223-3, a high copy-number plasmid, and transformed in E. coli. The presence of lacR in the resulting clone, termed pLL56, was verified by sequencing. Upon 2mM IPTG induction an expression of the gene could be obtained.

**Example 3****Cloning of the *L. delbrueckii* lac promoter in front of the chloramphenicol acetyltransferase gene in *Escherichia coli***

The lac promoters of *L. delbrueckii* LL44, N141 and N299 were tentatively cloned in front of an *E. coli* reporter gene, the chloramphenicol acetyltransferase (cat) gene. They were PCR amplified using the following primers

Eco RI		
ATATTAG <u>AATTC</u> AGTGACTTAAACTGG	SEQ ID No. 12	
Eco RI		
ATATTAG <u>AATTC</u> AGTACTTTGACACCG	SEQ ID No. 13	
Eco RI		
ATATTAG <u>AATTC</u> AAGAGGCTATATCGC	SEQ ID No. 14	
GGTTAATGCCGCCAAAGT	SEQ ID No. 15	

which contain an EcoRI site and primer 250 which is located in the lacS gene. The amplified fragments were digested with EcoRI and BspEI, which cleave in the primer sequence and 170 bp downstream the start codon of lacS. Restriction sites were filled with T<sub>4</sub> DNA polymerase, cloned in front of the promoterless cat gene of pKK232-8 digested with SmaI and transformed in *E. coli*. Chloramphenicol resistant clones were obtained for the three lactobacilli lac promoters, which indicates that the lac promoters of *L. delbrueckii* are active in *E. coli*. They were called pLL55 for N299, pLL57 for N141 and pLL58 for LL44 (Table 2) and all the constructs were verified by sequencing.

**Example 4****Control of the *L. delbrueckii* lac promoters by the LL44 lacR gene in *E. coli***

To test if the *L. delbrueckii* lac promoters could be controlled by the LL44 lacR in *E. coli*, plasmids pLL55, pLL57 and pLL58 were cotransformed with a low copy number plasmid containing the LL44 lacR gene under the control of the promoter of *E. coli* lac repressor (lacI). To obtain this construct, the two DNA fragments were linked by the Gene Splicing

by Overlap Extension (gene SOEing) method (Horton R., Molecular Biotechnology 3 (1995), 93 – 99).

The promoter of lacI was first amplified from pET11c using the following primers

Pst I	
ATAAATCTGCAGTGGGTATGGTGGC	SEQ ID No. 16
GATCGTTGCCACATTCACCACC	SEQ ID No. 17

The primer SEQ ID No. 18 is composed of a sequence of the lacI promoter and of the *L. delbrueckii* 5' end of lacR. The complete lacR gene was then PCR amplified using the following primers

GGTGAATGTGGCAACGATCAG	SEQ ID No. 18
Pst I	
ATATTACTGCAGACAGAATGCAGCC	SEQ ID No. 19

The sequence of primer SEQ ID No. 19 is the complement of SEQ ID No. 20. Both PCRs were purified, mixed and reamplified with primers SEQ ID No. 16 and SEQ ID No. 19, linking the lacI promoter and the LL44 lacR gene. Additionally, the ATG start codon of the lacR gene was replaced by GTG as in lacI (Fig. 6). The whole construct was cloned in the PstI site of the low copy number plasmid pACYC177 (New England Biolabs) digested with PstI and resulted after transformation in plasmid pLL62 (Fig. 6). The presence of the modified lacR gene in pLL62 was confirmed by digestions and PCR amplifications.

This plasmid was cotransformed with pLL55, pLL57 and pLL58 in *E. coli* resulting in strains LZL63, LZL64 and LZL65 (Table I) respectively. The resulting strains contained the different *L. delbrueckii* promoters together with the LL44 lacR gene. Regulation of the promoters was tested following the chloramphenicol acetyltransferase assay of Shaw, supra in the presence of lactose or glucose. The results showed that after 4 min incubation at 37°C, strains containing the different promoters were fully induced in YT medium or in a minimal medium based on M9-salts in the presence or in the absence of lacR as shown in the table below :

**Table I**  
chloramphenicol acetyltransferase assay (Shaw, 1975)

Plasmid	Sugar	YT-broth	Sugar	M9-salts
		OD <sub>415</sub>		OD <sub>415</sub>
pKK232-8	glucose	1.8		
pKK232-8	lactose	1.9		
LZL63	glucose	2.6	glucose	0.7
N299 prom. (pLL55) + lacR (pLL62)	lactose	2.5	lactose	0.9
LZL64	glucose	2.5	glucose	0.8
N141 prom. (pLL57) + lacR (pLL62)	lactose	0.4	lactose	1.2
LZL65	glucose	1.2	glucose	1.3
LL44 prom. (pLL58) + lacR (pLL62)	lactose	1.1	lactose	1.0

The OD<sub>415</sub> was read following 4 min incubation at 37°C.

The same results were found for the Gram-positive bacterium *Staphylococcus aureus*, where the lac promoter was functional in *E. coli* but the lacR gene not (Oskouian et al, J. Bacteriol. **169** (1990), 5459-5465). One possible explanation is that an additional factor, present only in Gram-positive bacteria, is necessary for proper repressor function.

### **Example 5**

#### **Cloning of the *L. delbrueckii* lac promoters in front of the $\beta$ -glucuronidase gene in *Lactococcus lactis***

The lac promoters of LL44 (inducible) and N299 (constitutive) were cloned in front of a reporter gene, the  $\beta$ -glucuronidase (*gusA*) gene. They were PCR amplified using primers SEQ ID No. 12 and No. 14, respectively, both containing an EcoRI site and primer SEQ ID No. 15 which is located in the lacS gene. The amplified fragments were digested with BspEI, located 170 bp downstream the lacS start codon, filled with T<sub>4</sub> DNA polymerase, and digested with EcoRI. Fragments were then cloned in front of the promoterless *gusA*



gene of pNZ272 (Platteeuw et al, Appl. Env. Microb. **60** (1994), 587-593) digested with Avall (refilled with T<sub>4</sub> DNA polymerase) and EcoRI. The resulting plasmids, pLL110 for LL44 and pLL113 for N299, were transformed in *Lc. lactis* (MG1363, plasmid free) (Fig. 7 and 8).

Chloramphenicol resistant clones were screened in the presence of 5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronide (X-glu), which is cleaved by the glucuronidase generating a blue color due to the liberation of the indol group of X-glu. Both plasmids were able to express  $\beta$ -glucuronidase, indicating that the lac promoter of *L. delbrueckii* is functional in *Lc. lactis*. The presence of the different promoters was checked by sequencing.

### **Example 6**

#### **Cloning of the *L. delbrueckii* lacR gene behind the $\beta$ -glucuronidase gene in *Lactococcus lactis***

To test if the lac promoters can be controlled by the *L. delbrueckii* lac repressor, the lacR gene of LL44 was cloned on plasmids pLL110 and pLL113. The lacR gene was PCR amplified using the following oligonucleotides:

Xho I  
ATAAATCTCGAGTGGTGATTGAGCC SEQ ID No. 20

Xho I  
ATATTACTCGAGACAGAATGCAGCC SEQ ID. No. 21

both containing a XhoI site. The primers were chosen to induce flanking regions of the gene. That is the oligonucleotide SEQ ID No. 20 is located 80 bp upstream the lacR start codon in the lacZ gene and the oligonucleotide SEQ ID No. 21 70 bp downstream the lacR stop codon, in the Asn-tRNA synthetase (asnA) gene. The fragment was cleaved with XhoI and cloned in the same site downstream the gusA gene of pLL110 and pLL113. With pLL110 containing the LL44 lac promoter, one clone (pLL112) was obtained with a lacR gene inserted in the opposite (rev) orientation as the gusA gene (Fig. 7). With pLL113 containing the N299 promoter, the lacR gene could be cloned in both orientations, pLL115 in the same (for) and pLL116 in the reverse (rev) orientation as the gusA gene (Fig. 8). The

mutated lac repressor ( $\otimes$ lacR) gene of LZL102 was PCR amplified and cloned as described above for LL44, resulting in plasmids pLL111 with the lac promoter of LL44 and pLL114 with the promoter of N299, both in the same orientation as gusA.

### **Example 7**

#### **Control of the $\beta$ -glucuronidase activity in *Lactococcus lactis***

The different constructs containing the *L. delbrueckii* lac promoters and repressors were studied by growing the transformed *Lc. lactis* in the presence of mannose. This sugar was shown to have no influence on the lactose metabolism of *Lactococcus*, as compared to saccharose and cellobiose. Exponentially growing cells were diluted into fresh media containing different concentrations of lactose and glucose, grown and harvested.  $\beta$ -glucuronidase (GusA) activity was measured from permeabilized cells in the presence of p-nitrophenol- $\beta$ -D-glucuronide, which is cleaved by GusA, liberating para-nitrophenol staining the solution in yellow. The results of these experiments are shown in figure 9.

### **Example 8**

#### **Regulation of the lac promoter of *L. delbrueckii* subsp. *lactis* LL44**

The promoter of LL44 (pLL110) was functional in *Lc. lactis* resulting in the production of  $\beta$ -glucuronidase in the presence of lactose or mannose. When glucose was added to the medium with or without lactose, a five to seven fold reduction in GusA activity was observed. This indicates that the glucose catabolite repression is efficient in the system.

**Table II**  
Expression of the gusA gene fused to the *L. delbrueckii* lac promoter  
in *Lactococcus lactis* MG1363 grown in M7

Beta-glucuronidase activity (U/mg prot. x 10 <sup>-4</sup> ) <sup>a</sup>									
Sugar added	<i>L. lactis</i> LL44 promoter				<i>L. bulgaricus</i> N299 promoter				
	pLL110	pLL111	pLL112	pLL113	pLL114	pLL115	pLL116		
	-	+⊗lacR <sup>b</sup>	+lacR	- ←	+⊗lacR	+lacR	+lacR	→	←
0.5% mannose	20.1±8.0	35.7±9.8	0.5±0.5	35.8±0.5	9.9±0.6	1.1±1.2	0.7±0.7		
0.05% lactose	15.6±4.4	35.8±7.1	3.4±1.1	32.0±4.1	10.5±2.2	0.7±0.7	0.7±0.8		
0.2% lactose	15.1±4.5	34.1±7.5	20.7±13.4	34.4±7.2	8.2±1.6	0.9±1.2	0.3±0.1		
0.5% lactose	20.1±12.6	37.0±7.8	27.1±7.5	34.1±9.4	8.5±1.1	1.5±1.0	11.1±1.3		
1.0% lactose	15.3±6.0	29.7±1.3	23.8±8.6	33.7±7.0	8.7±0.8	3.2±0.5	12.8±4.8		
0.5% glucose	5.1±3.7	34.6±9.1	0.5±0.2	22.3±2.0	5.6±1.0	0.7±0.8	0.6±0.6		
0.5% glucose +									
0.5% lactose	3.5±1.4	32.5±4.9	3.3±1.6	28.0±7.6	6.0±0.8	0.6±0.6	2.0±0.7		
0.5% glucose +									
1.0% lactose	3.7±0.9	38.7±12.0	4.8±2.2	22.5±6.6	4.7±1.0	0.6±0.2	4.0±1.2		

In the presence of the repressor gene (pLL112), GusA activities were similar to those observed with pLL110 when lactose (0.2 to 1%) was added. Almost no activity was observed with mannose (Table 2) indicating that the repressor is expressed in *Lc. lactis* and can bind to the lac promoter. Galactose, which is a common inducer of the lac system showed no induction of the system. Plasmid pLL112 was also subjected to the glucose catabolite repression, with a five to nine fold reduction in activity.

Surprising results were obtained with pLL111, the plasmid which contains the  $\odot$ lacR of LZL1012 and the LL44 lac promoter. GusA activities were twice higher than with the lac promoter alone even in the presence of glucose (Table II). The mutated repressor, with half of the protein translated, seems to play the role of an enhancer.

### **Example 9**

#### **Regulation of the lac promoter of *L. delbrueckii* subsp. *bulgaricus* N299**

The lac promoter of N299 (pLL113) induced production of GusA activity, almost twice as much as that produced by the promoter of LL44 (Table II, above). The N299 promoter is thus also functional in *Lc. lactis*. In the presence of glucose, only a relatively weak repression was observed, which could be explained by a different sequence of the CRE motif.

In the presence of the LL44 lacR gene, cloned in both orientations behind the gusA gene (pLL115, for and pLL116, rev), almost no GusA activity was detected in the presence of mannose. This result indicates that the so called "constitutive" promoter of *L. delbrueckii* subsp. *bulgaricus* is in fact repressed by the *L. delbrueckii* lac repressor. In addition, increasing concentrations of lactose induced increasing production of GusA activity. Nevertheless, the production even at 1% lactose with the repressor in the reverse orientation (pLL116) reached only half of that obtained in the absence of repressor. The N299 is not only regulated by the repressor, but more tightly than that of LL44 (Table V).

Finally in the presence of the mutated repressor (pLL114), the GusA activity was 3 to 4 times lower than in the absence of any repressor. On this promoter the  $\odot$ lacR seems to have a general inhibiting activity as a low activity is observed with or without lactose.

**Example 10****Lactose induction of *L. delbrueckii* promoters in *Lactococcus lactis***

The GusA activity was measured at several times during exponential growth of *Lc. lactis* containing the lac promoters of LL44 and N299 and the lacR gene of LL44 (pLL112, pLL115 and pLL116) in the presence of different concentrations of lactose and glucose (Fig. 9). The lac promoter of LL44 was maximally induced with 1% lactose and clearly subjected to glucose catabolite repression, i.e. 0.5% glucose in addition to lactose reduced this activity by half. As expected, mannose and glucose alone generated no GusA activity. Plasmids containing the N299 lac promoter (pLL115, for and pLL116, rev) are hardly inducible and are completely inhibited by glucose. Plasmid pLL115, having the repressor gene in the same orientation as gusA, resulted in the lowest lactose induction. In this case, it can be postulated that the repressor is produced in large amounts under the control of the *L. delbrueckii* lac promoter whereas in pLL116, it must be under the control of a weak promoter present on pNZ273 and is thus produced in smaller amounts.

**Example 11** **$\beta$ -galactosidase activity in *Lactobacillus delbrueckii***

The relation between the lac promoter and the lac repressor were studied in situ.  $\beta$ -galactosidase ( $\beta$ -gal) activity was measured in different *L. delbrueckii* strains in BHI-broth containing different sugars. The *L. delbrueckii* subsp. *lactis* strains tested showed no or low  $\beta$ -gal activity in the presence of mannose/galactose or glucose

**Table III**  
Expression of the lacZ gene in different strains of  
*Lactobacillus delbrueckii* grown in BHI

Beta-galactosidase activity (mU/OD600)							
Sugar added	LL44	LZL102	LB68	N62	N141	LB10	N299
0.5% galactose	2.4/1.8	6.4/9.5	2.4/2.0	0.0/0.0	0.6/0.5 <sup>a</sup>	15.3/12.6	7.9/14.9
0.02% lactose	14.1/17.5	12.1/9.5	21.2/20.6	11.5/9.5	13.5/9.7	18.0/12.1	14.3/10.6
0.05% lactose	11.4/13.9	7.4/9.5	24.3/20.6	11.6/9.1	11.0/10.9	16.9/8.6	9.5/18.1
0.2% lactose	10.1/12.3	11.6/9.5	14.1/20.3	8.8/8.1	7.6/8.3	10.2/7.4	23.8/23.8
0.5% lactose	19.0/8.7	10.6/9.5	20.4/16.5	15.4/12.4	9.8/5.5	12.8/11.1	29.0
0.5% glucose	0.9/0.6	14.7/9.5	3.2/1.9	0.0/0.0	2.5/2.2	12.6/7.8	39.7/15.5
0.5% glucose + 0.5% lactose	15.9/7.1	11.7/9.5	15.6/16.8	17.9/33.1	6.3/5.4	8.0/7.6	28.6/31.7
0.5% glucose + 1.0% lactose	7.7/6.7	16.6/9.5	15.4/15.5	14.9/13.1	9.3/7.0	12.0/10.3	22.0/23.8

a : galactose was replaced by mannose

At all concentrations of lactose tested (0.02 to 0.5%), the  $\beta$ -gal activity was in the same range, indicating that very low concentration of lactose is able to fully induce the expression of the lac operon. In the presence of lactose and glucose, strains were not subjected to catabolite repression and even a stimulation of  $\beta$ -gal activity was found for N62 (Table 3, above). The constitutive *L. delbrueckii* subsp. *bulgaricus* N299 showed the same  $\beta$ -gal activity with all sugars used even in the presence of glucose. Two mutants of *L. delbrueckii* subsp. *lactis*, LB10 and LZL102, were also analysed. LZL102, the spontaneous mutant of LL44, produces a truncated repressor and LB10 produces no peptide at all (Fig. 12). In both cases, strains were constitutive with equal amounts of  $\beta$ -gal produced with galactose, lactose or glucose.

### **Example 12**

#### **Lactose induction of the lac promoters in *Lactobacillus delbrueckii***

The  $\beta$ -gal induction was measured at several times during exponential growth of the *L. delbrueckii* subsp. *lactis* strain LL44 and the *bulgaricus* strain N299. The maximum level of activity was reached in both strains after 15 min incubation (Fig. 10). For LL44, no  $\beta$ -gal activity was detected with galactose or glucose, whereas full induction was obtained with all lactose concentrations. The enzymatic activity was slightly reduced in the presence of glucose, i.e. a full activity was only reached after 60 min. For N299, all sugars resulted in the same induction with no glucose catabolite repression.

### Claims

1. A DNA sequence represented by the following general formula

$$\begin{aligned} & \text{p/o} - (\text{A})_n - \text{R}_y, \text{ or} \\ & \text{p/o} - \text{R}_y - (\text{A})_n \end{aligned}$$

wherein

p/o denotes the DNA sequence identified under SEQ ID No. 9 or a functional variant thereof, which retains its capability to bind to the lac repressor protein of *Lactobacillus delbrueckii*;

A denotes a gene coding for a polypeptide of interest,

n denotes an integer of  $\geq 0$ ;

R denotes a gene coding for the lac repressor protein as identified under SEQ ID No. 2 or a functional variant thereof; and

Y is 0 or 1.

2. The DNA sequence according to claim 1, wherein y is 1.
3. The DNA sequence according to claim 1, wherein the reading frame of gene coding for the lac repressor is reversed relative to the region p/o.
4. The DNA sequence according to claim 1 wherein the gene coding for a polypeptide of interest is selected from group consisting of genes encoding enzymes, cell surface proteins, or functional peptides.
5. The DNA sequence according to claim 4, wherein the gene coding for a polypeptide of interest is selected from the group consisting of genes coding for dextranucrase, glycosyltransferase, phytase, transglutaminase, peptidase, phenylalanine ammonia lyase, protease, cell surface antigens, bacteriocins, hormones or insulin.



6. The DNA sequence according to any of the preceding claims, which is devoid of catabolite responsive elements.
7. A DNA sequence coding for the lac repressor protein of *lactobacillus delbrueckii* as identified by SEQ ID No. 2 or a functional variant thereof retaining the capability to bind to the DNA sequence as identified by SEQ ID No. 9.
8. A recombinant microorganism harboring a DNA sequence according to any of the preceding claims.
9. The microorganism according to claim 8, which is a gram positive bacterium.
10. The microorganism according to claim 8 or 9, which is selected from the group consisting of lactic acid bacteria.
11. The microorganism according to any of the claims 8 to 10, wherein the DNA sequence of claims 1 to 7 is incorporated into the bacteria's chromosome or is harbored in a plasmid maintained extra-chromosomal.
12. The microorganism according to claim 8, which is CNCM I-2089, CNCM I-2090 or CNCM I-2091.
13. Use of a DNA sequence according to any of the claims 1 to 7 for the production of a polypeptide A.
14. The use according to claim 13, wherein the DNA sequence is harbored in a plasmid maintained extra-chromosomal or is in the bacteria's chromosome.
15. The use according to any of the claims 13 or 14, wherein expression is performed in gram positive microorganisms.

16. The use according to any of the claims 13 to 15, wherein expression is performed in microorganisms selected from the group consisting of lactic acid bacteria.
17. The use of a microorganism according to any of the claims 8 to 11 for the production of food products.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau




(43) International Publication Date  
11 January 2001 (11.01.2001)

PCT

(10) International Publication Number  
WO 01/02576 A1

- (51) International Patent Classification<sup>7</sup>: C12N 15/31, 15/74, C07K 14/335, A23C 9/123 43, chemin d'En-Quettolla, CH-1616 Attalens (CH).  
MOLLET, Beat [CH/CH]; 9, avenue de l'Esplanade, CH-1012 Lausanne (CH).
- (21) International Application Number: PCT/EP00/05834
- (22) International Filing Date: 23 June 2000 (23.06.2000) (74) Agent: STRAUS, Alexander; Becker, Kurig, Straus, Bavariastrasse 7, D-80336 München (DE).
- (25) Filing Language: English (81) Designated States (national): AU, BR, CA, JP, US.
- (26) Publication Language: English (84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
- (30) Priority Data: 99112471.0 30 June 1999 (30.06.1999) EP
- (71) Applicant (for all designated States except US): SOCI-  
ETE DES PRODUITS NESTLE S.A. [CH/CH]; P.O. Box 353, CH-1800 Vevey (CH).  
30 Dec 01/30 mos Published:  
With international search report.  
Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- (72) Inventors; and
- (73) Inventors/Applicants (for US only): GERMOND, Jacques, Edouard [CH/CH]; 18, chemin des Cèdres, CH-1023 Crissier (CH). LAPIERRE, Luciane [CH/CH];  
For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: THE LACTOSE OPERON OF LACTOBACILLUS DELBRUECKII AND ITS USE FOR CONTROLLING GENE TRANSCRIPTION AND/OR EXPRESSION IN BACTERIAL CELLS



Physical map of the lactose operon of the different *L. delbrueckii* studied. Open arrows are for the lac operon genes and dashed arrow is for inactivated lacR. Boxes are for the different IS-elements, where the arrows heads are for the inverted repeats.  
\* same sequence as LL44 except an insertion in the 5' end of the lacA gene.

(57) Abstract: The present invention relates to a DNA sequence suitable for the controlled transcription and/or expression of a variety of different genes in bacteria, preferably gram positive bacteria. In particular the present invention pertains to a DNA sequence comprising the promoter and the gene coding for the lac repressor of the lac operon of *Lactobacillus delbrueckii* with a DNA sequence coding for a gene product of interest being arranged inbetween.

WO 01/02576 A1

01




Fig. 2




Fig. 3



Fig. 2 u. 3 : Organisation of the promoter region of LL44 and N299 *lac* operon. Operators O1 and O2 are indicated by black boxes. The inverted repeats of the operators are represented by arrows. The sequence responsible for catabolite repression (CRE) is overdrawn by stars. The inverted repeat of ISL5 is boxed and shaded. The initiation of transcription is shown by an i (arrow head) (Leong-Morgenthaler et al, 1991). The promoter sequence of LL44 is numbered according to figure 1. The picture is not drawn to scale.

```

1   GAATTTTGTCTGGATGCTCAGGAAGCCCGCCAGCTCAAGCTGGTGATTAGCCACTTTTT
    stop lacZ
61  ACTGAATATGCTACAATTGACTTAACAGCATAAAATTTTAGTAAAAGCGAGTGAAGAAG
    RBS
121  ATGGCAACGATCAGAGAAGTGGCCAAGGCAGCCGGCGTGTGCCAGCGACGGTTTCCCGG
1   M A T I R E V A K A A G V S P A T V S R
    helix          turn          helix
181  GTCTTGAAGTATGACCAGACCCTGTGGTCAATGAGGCAACGCGGCAGAAGATATTCAA
21  V L N Y D Q T L S V N E A T R Q K I F K
241  ACTGCTGAAGCCATGCACTACCATAAGAGCCGGAAGACCAGAAAGAGCAAGCAAAAGCGC
41  T A E A M H Y H K S R K T R K S K Q K R
301  CTGGCGATCTGCCTGTGGTGTGACCAAGACCAGGAGATCAAGGACCTCTATTACTATTCA
61  L A I C L W C D Q D Q E I K D L Y Y Y S
361  ATCAGAACCAGCGCGCAAGCAGAGGCCAAGAAGCAGGGACTTGAAAGCCAGGTCAATTTAT
81  I R T S A Q A E A K K Q G L E S Q V I Y
421  CCGGCTGATCCTTTGCCGATCCAGCTGCTTTAAGCGGGATTATCATGATTGGCTACCAG
101  P A D P L P D P A A L S G I I M I G Y Q
481  CAGTATTCGCCAGACCGCTTGAATGAAGTCAAAAAGTCTGGCCTGCCCCTGGTCTTTGTC
121  Q Y S P D R L N E V K K S G L P L V F V
541  GATACTGACACCTTAAATTTGGGTTACTGCTCAGTTGTGGCTGACTTTGGCCAGGCCATG
141  D T D T L K L G Y C S V V A D F G Q A M
601  CAGGAGGCGCTAGAGGTCTTCTGGGGGCAGGGCAGGGAGCGGATCGCCCTTTGGATGGT
161  Q E A L E V F W G Q G R E R I A L L D G
661  GATTTGGACAGTAATTTTGATAAAAACAACCTTGGTCGACTTCCGCTTCCGCGATTATAAG
181  D L D S N F D K N N L V D F R F R D Y K
    ▼
721  AAGAGCCTCGCGGCCCGCGGCCAGTACGACCCGGACTTAGTCTATGTTGGAACTTCACT
201  K S L A A R G Q Y D P D L V Y V G N F T
781  CCGCAATCTGGCTATGAAGCCATTAAAGAAGCTCTTAAGTCCGGCTCCTTCCCGAAAGCC
221  P Q S G Y E A I K E A L K S G S F P K A
841  TTGATTGCGGCTAATGACGCCATGGCTATTGGAGCATTGAAGGCCTTTAAAGAAGCTGGA
241  L I A A N D A M A I G A L K A F K E A G
901  ATTAAGTCCCAGAGGACGTGAGTCTGATTTCTTTAATGACACAACGGCAGCAGAATTT
261  I K V P E D V S L I S F N D T T A A E F
961  GCCAACCCAGCCTTGACTAGCGTACATGTAGAGAC(CAG)CAGATGGGCCGAGCCAGCGTC
281  A N P A L T S V H V E T Q Q M G R A S V
1021 AAGGTCATGAAAGACCTGCTGGATGATGATGAAGCCGGCACTTACAAGGTCACCTTTCCCA
301  K V M K D L L D D D E A G T Y K V T F P
1081 ACAAACCTCGTTTACCGGAATCTTGCCAAAAGCATAAGGGCATAGAGCATAATAACAG
321  T K L V Y R E S C P K A *
    →
1141 CAAAGAAATAGCTTGAGATTGATTTCTCCAAGCTATTTTCGTATATATTGCGTGC
    stop asnA
1201 ATTCTGTTGATCATTCTTGGGAATGGGACAGCTTCACGAACGTGGTCCAGCTTGAGATC
1261 CAGGCAATGACCCGTTCAAAG

```


Figure 4:

Nucleotide and amino acid sequences of the *L. delbrueckii* subsp. *lactis* LL44 *lacR* gene. Start (121) and stop (1119) codons are boxed. Putative *lacR* RBS is underlined. The putative rho-independent terminator is underlined by convergent arrows. Stop codons of the beta-galactosidase (*lacZ*) and Asn t-RNA synthetase (*asnA*) genes are boxed. Insertion sequence of ISL3 is represented by a large open arrow. Single base pair deletion (722) in the mutant LZL102 is shown by an arrow head, leading to a premature stop codon (758) underlined.




Physical map of the lactose operon of the different *L. delbrueckii* studied. Open arrows are for the *lac* operon genes and dashed arrow is for inactivated *lacR*. Boxes are for the different IS-elements, where the arrows heads are for the inverted repeats.

- same sequence as LL44 except an insertion in the 5' end of the *lacA* gene.




**Figure 6 :** Schematic representation of the construction of a pLL62. The dark box is for LL44 *lacR* gene and the white box is for the promoter region of the *lacI* gene of pET11c. Both were linked by PCR amplification using the SOEing method.







**Figure 7** : Schematic representation of the construction of pLL110 and pLL112. Dashed arrows are for the genes of the *L. delbrueckii* lac operon, and open arrows for plasmid genes. The dark box is for the promoter region cloned in front of the *gusA* gene. Plasmids are not drawn to scale. The simple arrows represent the primers used to amplify the cloned regions.



**Figure 8** : Schematic representation of the construction of pLL113, pLL115 and pLL116. Dashed arrows are for the genes of the *L. delbrueckii* *lac* operon, and open arrows for plasmid genes. The open box containing arrow heads represents the IS-elements. The dark box is for the promoter region cloned in front of the *gusA* gene. Plasmids are not drawn to scale. The simple arrows represent the primers used to amplify the cloned regions.



**Figure 9:** Beta-glucuronidase activity (mean of three experiments) of *Lactococcus lactis* MG1363 containing different *Lactobacillus delbrueckii* *lac* promoter and the *lacR* gene of LL44. The *lacR* orientation compared to the *gusA* gene is represented by an arrow. The medium used was M17 containing : 0.5% mannose(\*), 0.05% lactose (▲), 0.2% lactose (●), 0.5% lactose (■), 1.0% lactose (◆), 0.5% glucose (△), 0.5% glucose + 0.5% lactose (□) and 0.5% glucose + 1.0% lactose (◇).



**Figure 10:** Beta-galactosidase activity (mean of three experiments) of *Lactobacillus delbrueckii* subsp. *lactis* LL44 and *L. delbrueckii* subsp. *bulgaricus* N299. The medium used was BHI-broth containing : 0.5% galactose (✕), 0.05% lactose (▲), 0.2% lactose (●), 0.5% lactose (■), 1.0% lactose (◆), 0.5% glucose (△), 0.5% glucose + 0.5% lactose (◻), and 0.5% glucose + 1.0% lactose (◊). Strain N299 did not grow neither in galactose alone nor in 0.05% lactose and the experiment was not realised with these sugars.

# DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

LACTOSE OPERON OF LACTOBACILLUS DELBRUECKII AND ITS USE FOR CONTROLLING GENE TRANSCRIPTION AND/OR EXPRESSION IN BACTERIAL CELLS

the specification of which: (check one)

- ☐ is attached hereto.
- ☒ was filed on 23 June 2000, as United States Application No. or PCT International Application No. PCT/EP00/05834 and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent Office all information which is known to me to be material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code Section 119 or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT international application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Number	Country	Day/Month/Year Filed	Priority Not Claimed
99112471.0	Europe	30 June 1999	<input type="checkbox"/>
			<input type="checkbox"/>
			<input type="checkbox"/>

Docket No. 112843-039

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

**Application Serial No.**

**Filing Date**

I hereby claim the benefit under 35 U.S.C. Section 120 of any United States application(s), or Section 365(c) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C.F.R., Section 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

**Application Serial No.**

**Filing Date**

**Status**  
**(patented, pending, abandoned)**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.


And I hereby appoint the practitioners at customer number: 29174





as my attorneys with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and direct that all correspondence be forwarded to:

Bell, Boyd & Lloyd LLC  
P.O. Box 1135  
Chicago, Illinois 60690-1135

And all telephone calls be directed to: (312) 807-4204.

Full name of sole or first inventor	<b>Jacques Edouard Germond</b>	
Sole or first inventor's signature		Date 18.02.02
Residence	CH-1023 Crissier, Switzerland	
Citizenship	Switzerland	
Post Office Address	18, chemin des Cèdres	
	CH-1023 Crissier, Switzerland	

Full name of second inventor <b>Luciane LaPierre</b>	
Second inventor's signature <i>X d. Lapierre</i>	Date 23.02.02
Residence CH-1616 Attalens, Switzerland <i>CHX</i>	
Citizenship Switzerland	
Post Office Address 43, chemin d'En-Quettolla	
CH-1616 Attalens, Switzerland	

Full name of third inventor <b>Beat Mollet</b>	
Third inventor's signature 	Date 1/18/2002
Residence CH-1012 <u>Lausanne</u> , Switzerland 	
Citizenship Switzerland	
Post Office Address 9, avenue de l'Esplanade	
CH-1012 Lausanne, Switzerland	

10/01/98 13:13:17  
Rec'd PCT/PTO 13 MAY 2004  
10/01 9817

SEQUENCE LISTING

<110> Société des Produits Nestlé S.A.

<120> The Lactose operon of Lactobacillus delbrueckii and its  
use for controlling gene transcription and/or  
expression in bacterial cells

<130> 112843-039

<140> PCT/EP00/05834

<141> 2000-06-23

<150> 99112471.0

<151> 1999-06-30

<160> 22

<170> PatentIn Ver. 2.1

<210> 1

<211> 1435

<212> DNA

<213> Lactobacillus delbrueckii

<400> 1

```
gaattttgtc tggatgctca ggaagccgc cagctcaagc tggtgattca gccacttttt 60
actgaataat gctacaattg acttaacagc ataaaaatttt agtaaaagcg agtgaagaag 120
atggcaacga tcagagaagt ggccaaggca gccggcgtgt cgccagcgac ggtttcccg 180
gtcttgaact atgaccagac cctgtcggtc aatgaggcaa cgcggcagaa gatattcaaa 240
actgctgaag ccattgacta ccataagagc cggaagacca gaaagagcaa gcaaaagcgc 300
ctggcgatct gcctgtggtg tgaccaagac caggagatca aggacctcta ttactattca 360
atcagaacca gcgcgcaagc agaggccaag aagcagggac ttgaaagcca ggtcatttat 420
ccggctgac ctttgcccga tccagctgct ttaagcggga ttatcatgat tggctaccag 480
cagtattcgc cagaccgctt gaatgaagtc aaaaagtctg gcctgcccct ggtctttgtc 540
gatactgaca ccttaaaatt gggttactgc tcagttgtgg ctgacttttg ccaggccatg 600
caggaggcgc tagaggtctt ctgggggcag ggcagggagc ggatcgccct tttggatggt 660
gatttggaca gtaattttga taaaaacaac ttggtcgact tccgcttccg cgattataag 720
aagagcctcg cggcccgcgg ccagtagcac ccggacttag tctatgttgg aaacttcact 780
ccgcaatctg gctatgaagc cattaagaa gctcttaagt ccggtcctt cccgaaagcc 840
ttgattgcgg ctaatgacgc catggctatt ggagcattga aggcctttaa agaagctgga 900
attaaagtcc cagaggacgt cagtctgatt tcttttaatg acacaacggc agcagaattt 960
gccaacccag ccttgactag cgtacatgta gagaccagc agatgggccc agccagcgctc 1020
aaggtcatga aagacctgct ggatgatgat gaagccggca cttacaaggc cactttccca 1080
acaaaactcg tttaccggga atcttgcca aaagcataag ggcatagagc ataataacag 1140
caaagaaata gcttggagat tgattttctc caagctatct ttcgtatata ttatggctgc 1200
attctgttga tcattcttgg gaatgggaca gcttcacgaa cgtgggtccag cttgcagatc 1260
caggcaatga cccgttcaaa gcccatcccg aagccggagt gcggcacgtg ccgtactttt 1320
ctcagggtcc caggtagcca ggagtagtcg tcccagggtt gagggccgct tcttcgattt 1380
gcgccttcaa ggtgtcgtag tcagcttcac gttctgatcc gccatgattt cccgt 1435
```

<210> 2

<211> 332

<212> PRT

<213> Lactobacillus delbrueckii



<400> 2

Met 1	Ala	Thr	Ile	Arg 5	Glu	Val	Ala	Lys	Ala	Gly	Val	Ser	Pro	Ala 15	
Thr	Val	Ser	Arg 20	Val	Leu	Asn	Tyr	Asp 25	Gln	Thr	Leu	Ser	Val 30	Asn	Glu
Ala	Thr	Arg 35	Gln	Lys	Ile	Phe	Lys 40	Thr	Ala	Glu	Ala	Met 45	His	Tyr	His
Lys	Ser 50	Arg	Lys	Thr	Arg	Lys 55	Ser	Lys	Gln	Lys	Arg 60	Leu	Ala	Ile	Cys
Leu 65	Trp	Cys	Asp	Gln	Asp 70	Gln	Glu	Ile	Lys	Asp 75	Leu	Tyr	Tyr	Tyr	Ser 80
Ile	Arg	Thr	Ser	Ala 85	Gln	Ala	Glu	Ala	Lys 90	Lys	Gln	Gly	Leu	Glu 95	Ser
Gln	Val	Ile	Tyr 100	Pro	Ala	Asp	Pro	Leu 105	Pro	Asp	Pro	Ala	Ala 110	Leu	Ser
Gly	Ile	Ile 115	Met	Ile	Gly	Tyr	Gln 120	Gln	Tyr	Ser	Pro	Asp 125	Arg	Leu	Asn
Glu 130	Val	Lys	Lys	Ser	Gly	Leu 135	Pro	Leu	Val	Phe	Val 140	Asp	Thr	Asp	Thr
Leu 145	Lys	Leu	Gly	Tyr	Cys 150	Ser	Val	Val	Ala	Asp 155	Phe	Gly	Gln	Ala	Met 160
Gln	Glu	Ala	Leu	Glu 165	Val	Phe	Trp	Gly	Gln 170	Gly	Arg	Glu	Arg	Ile 175	Ala
Leu	Leu	Asp 180	Gly	Asp	Leu	Asp	Ser	Asn 185	Phe	Asp	Lys	Asn 190	Asn	Leu	Val
Asp	Phe	Arg 195	Phe	Arg	Asp	Tyr	Lys 200	Lys	Ser	Leu	Ala	Ala 205	Arg	Gly	Gln
Tyr 210	Asp	Pro	Asp	Leu	Val	Tyr 215	Val	Gly	Asn	Phe	Thr 220	Pro	Gln	Ser	Gly
Tyr 225	Glu	Ala	Ile	Lys	Glu 230	Ala	Leu	Lys	Ser	Gly 235	Ser	Phe	Pro	Lys	Ala 240
Leu	Ile	Ala	Ala	Asn 245	Asp	Ala	Met	Ala	Ile 250	Gly	Ala	Leu	Lys	Ala 255	Phe
Lys	Glu	Ala	Gly 260	Ile	Lys	Val	Pro	Glu 265	Asp	Val	Ser	Leu	Ile 270	Ser	Phe
Asn	Asp	Thr 275	Thr	Ala	Ala	Glu	Phe 280	Ala	Asn	Pro	Ala	Leu 285	Thr	Ser	Val
His 290	Val	Glu	Thr	Gln	Gln	Met 295	Gly	Arg	Ala	Ser	Val 300	Lys	Val	Met	Lys

[illegible]

```
<210> 3'  
<211> 6  
<212> DNA  
<213> Lactobacillus delbrueckii
```

6

7

7

17

20

3

<400> 8  
tgtaagcgta aacaa 15

<210> 9  
<211> 98  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 9  
tgtttactaa aaatattttg gtaaagcatc ttgatttggt tagtaaacgg gtctatactg 60  
taagcgtaaa caagtttagaa cacctaaagg agaaaatc 98

<210> 10  
<211> 25  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 10  
atattactgc agagtaaaag cgagt 25

<210> 11  
<211> 25  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 11  
ataaataagc ttacagaatg cagcc 25

<210> 12  
<211> 27  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 12  
atattagaat tcagtgactt aaactgg 27

<210> 13  
<211> 27  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 13  
atattagaat tcagtacttt gacaccg 27

<210> 14  
<211> 27  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 14

atattagaat tcaagaggct atatcgc 27

<210> 15  
<211> 18  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 15  
ggttaatgcc gccaaagt 18

<210> 16  
<211> 25  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 16  
ataaatctgc agtgggtatg gtggc 25

<210> 17  
<211> 22  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 17  
gategttgcc acattcacca cc 22

<210> 18  
<211> 21  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 18  
ggtgaatgtg gcaacgatca g 21

<210> 19  
<211> 25  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 19  
atattactgc agacagaatg cagcc 25

<210> 20  
<211> 25  
<212> DNA  
<213> Lactobacillus delbrueckii

<400> 20  
ataaatctcg agtgggtgatt cagcc 25

<210> 21  
 <211> 25  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 21  
 atattactcg agacagaatg cagcc 25

<210> 22  
 <211> 98  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 22  
 tgtttactaa aaatattttg gttaaagcatc ttgatttggt tagtaaacgg gtctatactg 60  
 taagcgtaaa caagttagaa cacctaaagg agaaaatc 98

## SEQUENCE LISTING

&lt;110&gt; Societé des Produits Nestlé

&lt;120&gt; Lac-operon

&lt;130&gt; 80197

&lt;140&gt;

&lt;141&gt;

&lt;150&gt; 99112471.0

&lt;151&gt; 1999-06-30

&lt;160&gt; 22

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 1435

&lt;212&gt; DNA

&lt;213&gt; Lactobacillus delbrueckii

&lt;400&gt; 1

```

gaattttgtc tggatgctca ggaagcccg cagctcaagc tggtgattca gccacttttt 60
actgaataat gctacaattg acttaacagc ataaaatttt agtaaaagcg agtgaagaag 120
atggcaacga tcagagaagt ggccaaggca gccggcggtg cgccagcgac ggtttcccg 180
gtcttgaaact atgaccagac cctgtcggtc aatgaggcaa cgcggcagaa gatattcaaa 240
actgctgaag ccatgcacta ccataagagc cggaagacca gaaagagcaa gcaaaagcgc 300
ctggcgatct gcctgtggtg tgaccaagac caggagatca aggacctcta ttactattca 360
atcagaacca gcgcgcaagc agaggccaag aagcagggac ttgaaagcca ggtcatttat 420
ccggctgata ctttgccga tccagctgct ttaagcggga ttatcatgat tggctaccag 480
cagtattcgc cagaccgctt gaatgaagtc aaaaagtctg gcctgccctt ggtctttgtc 540
gatactgaca ccttaaaatt gggttactgc tcagttgtgg ctgacttttg ccaggccatg 600
caggaggcgc tagaggctct ctgggggcag ggcagggagc ggatcgccct tttggatgg 660
gatttggaac gtaattttga taaaaacaac ttggtcgact tccgcttcgc cgattataag 720
aagagcctcg cggcccgcg ccagtagcgc ccggacttag tctatggttg aaacttcact 780
ccgcaatctg gctatgaagc cattaaagaa gctcttaagt ccggctcctt cccgaaagcc 840
ttgattgcgg ctaatgacgc catggctatt ggagcattga aggcctttaa agaagctgga 900
attaaagtcc cagaggacgt cagtctgatt tcttttaatg acacaacggc agcagaattt 960
gccaaccag ccttgactag cgtacatgta gagaccagc agatgggccc agccagcgct 1020
aaggtcatga aagacctgct ggatgatgat gaagccggca cttacaaggt cactttccca 1080
acaaaactcg ttaccggga atcttgccca aaagcataag ggcatagagc ataataacag 1140
caaagaaata gcttgagat tgattttctc caagctattt ttcgtatata ttatggctgc 1200
attctgttga tcattcttgg gaatgggaca gcttcacgaa cgtgggtccag cttgcagatc 1260
caggcaatga cccgttcaa gcccatcccg aagccggagt gcggcacgtg ccgtactttt 1320
ctcagggtcc caggtaccca ggagtagtcg tcccagggtt gaggcccgct tcttcgattt 1380
gcgccttcaa ggtgtcgtag tcagcttcac gttctgatcc gccatgattt cccgt 1435

```

WO 01/02576

PCT/EP00/05834

<210> 2  
 <211> 332  
 <212> PRT  
 <213> Lactobacillus delbrueckii

<400> 2

Met Ala Thr Ile Arg Glu Val Ala Lys Ala Ala Gly Val Ser Pro Ala  
 1 5 10 15  
 Thr Val Ser Arg Val Leu Asn Tyr Asp Gln Thr Leu Ser Val Asn Glu  
 20 25 30  
 Ala Thr Arg Gln Lys Ile Phe Lys Thr Ala Glu Ala Met His Tyr His  
 35 40 45  
 Lys Ser Arg Lys Thr Arg Lys Ser Lys Gln Lys Arg Leu Ala Ile Cys  
 50 55 60  
 Leu Trp Cys Asp Gln Asp Gln Glu Ile Lys Asp Leu Tyr Tyr Tyr Ser  
 65 70 75 80  
 Ile Arg Thr Ser Ala Gln Ala Glu Ala Lys Lys Gln Gly Leu Glu Ser  
 85 90 95  
 Gln Val Ile Tyr Pro Ala Asp Pro Leu Pro Asp Pro Ala Ala Leu Ser  
 100 105 110  
 Gly Ile Ile Met Ile Gly Tyr Gln Gln Tyr Ser Pro Asp Arg Leu Asn  
 115 120 125  
 Glu Val Lys Lys Ser Gly Leu Pro Leu Val Phe Val Asp Thr Asp Thr  
 130 135 140  
 Leu Lys Leu Gly Tyr Cys Ser Val Val Ala Asp Phe Gly Gln Ala Met  
 145 150 155 160  
 Gln Glu Ala Leu Glu Val Phe Trp Gly Gln Gly Arg Glu Arg Ile Ala  
 165 170 175  
 Leu Leu Asp Gly Asp Leu Asp Ser Asn Phe Asp Lys Asn Asn Leu Val  
 180 185 190  
 Asp Phe Arg Phe Arg Asp Tyr Lys Lys Ser Leu Ala Ala Arg Gly Gln  
 195 200 205  
 Tyr Asp Pro Asp Leu Val Tyr Val Gly Asn Phe Thr Pro Gln Ser Gly

WO 01/02576

PCT/EP00/05834

210	215	220
Tyr Glu Ala Ile Lys Glu Ala Leu Lys Ser Gly Ser Phe Pro Lys Ala		
225	230	235 240
Leu Ile Ala Ala Asn Asp Ala Met Ala Ile Gly Ala Leu Lys Ala Phe		
	245	250 255
Lys Glu Ala Gly Ile Lys Val Pro Glu Asp Val Ser Leu Ile Ser Phe		
	260	265 270
Asn Asp Thr Thr Ala Ala Glu Phe Ala Asn Pro Ala Leu Thr Ser Val		
	275	280 285
His Val Glu Thr Gln Gln Met Gly Arg Ala Ser Val Lys Val Met Lys		
	290	295 300
Asp Leu Leu Asp Asp Asp Glu Ala Gly Thr Tyr Lys Val Thr Phe Pro		
305	310	315 320
Thr Lys Leu Val Tyr Arg Glu Ser Cys Pro Lys Ala		
	325	330

<210> 3  
 <211> 6  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 3  
 tgttta

6

<210> 4  
 <211> 7  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 4  
 gtaaaca

7

<210> 5  
 <211> 7  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 5



WO 01/02576

PCT/EP00/05834

gtaaacg

7

<210> 6

<211> 17

<212> DNA

<213> Lactobacillus delbrueckii

<400> 6

cgcttggtga ttcagcc

17

<210> 7

<211> 20

<212> DNA

<213> Lactobacillus delbrueckii

<400> 7

agctttacgg ggaagtcggg

20

<210> 8

<211> 15

<212> DNA

<213> Lactobacillus delbrueckii

<400> 8

tgtaagcgta aacaa

15

<210> 9

<211> 98

<212> DNA

<213> Lactobacillus delbrueckii

<400> 9

tgtttactaa aaatattttg gtaaagcatc ttgatttggt tagtaaacgg gtctatactg 60  
taagcgtaaa caagttagaa cacctaaagg agaaaatc 98

<210> 10

<211> 25

<212> DNA

<213> Lactobacillus delbrueckii

<400> 10

atattactgc agagtaaaag cgagt

25

WO 01/02576

PCT/EP00/05834

<210> 11  
 <211> 25  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 11  
 ataaataagc ttacagaatg cagcc

25

<210> 12  
 <211> 27  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 12  
 atattagaat tcagtgactt aaactgg

27

<210> 13  
 <211> 27  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 13  
 atattagaat tcagtacttt gacaccg

27

<210> 14  
 <211> 27  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 14  
 atattagaat tcaagaggct atatcgc

27

<210> 15  
 <211> 18  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 15  
 ggttaatgcc gccaaagt

18

<210> 16  
 <211> 25

WO 01/02576

PCT/EP00/05834

<212> DNA  
 <213> Lactobacillus delbrueckii

<400> 16  
 ataaatctgc agtgggtatg gtggc 25

<210> 17  
 <211> 22  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 17  
 gatcgttgcc acattcacca cc 22

<210> 18  
 <211> 21  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 18  
 ggtgaatgtg gcaacgatca g 21

<210> 19  
 <211> 25  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 19  
 atattactgc agacagaatg cagcc 25

<210> 20  
 <211> 25  
 <212> DNA  
 <213> Lactobacillus delbrueckii

<400> 20  
 ataaatctcg agtgggtgatt cagcc 25

<210> 21  
 <211> 25  
 <212> DNA  
 <213> Lactobacillus delbrueckii

WO 01/02576

PCT/EP00/05834

<400> 21

atattactcg agacagaatg cagcc

25

<210> 22

<211> 98

<212> DNA

<213> Lactobacillus delbrueckii

<400> 22

tgtttactaa aaatattttg gtaaagcatc ttgatttggt tagtaaacgg gtctatactg 60  
taagcgtaaa caagttagaa cacctaaagg agaaaatc 98